	<p>Policies and Procedures</p> <p>Title: PARENTERAL NUTRITION (PN) – ADULT ADMINISTRATION AND MAINTENANCE</p> <p>LPN Additional Competency (LPNAC): Parenteral Nutrition -Adult Administration and Maintenance with an Established Plan of Care RN Entry Level Competency</p> <p>I.D. Number: 1078</p>
<p>Authorization</p> <p>[x] Former SktnHR Nursing Practice Committee</p>	<p>Source: Nursing Date Revised: May 2018 Date Effective: January/February 2000 Scope: Former SktnHR Urban Acute Care and Home Care</p>

Any PRINTED version of this document is only accurate up to the date of printing 13-Aug-18. Former Saskatoon Health Region (SktnHR) cannot guarantee the currency or accuracy of any printed policy. Always refer to the Policies and Procedures site for the most current versions of documents in effect. Former SktnHR accepts no responsibility for use of this material by any person or organization not associated with former SktnHR. No part of this document may be reproduced in any form for publication without permission of the former SktnHR.

DEFINITIONS:

2 – in – 1 PN - consists of dextrose, amino acids, electrolytes, vitamins, minerals and trace elements with intravenous fat emulsion administered on separate lines.

3 – in – 1 PN – (Total Nutrient Admixture) consists of dextrose, amino acids, intravenous fat emulsion, electrolytes, vitamins, minerals and trace elements.

Central – formulation appropriate for delivery via a central line.

Client – Term used to refer to residents, patients and clients.

Creaming – A dense white colour that appears at the top of the formulation.

Established Plan of Care – based on RN assessment of care needs, the plan of care for PN Administration and Maintenance may be considered established for clients who have had PN running for at least two days without complications or for whom PN orders are written weekly. Plan of care must be documented in the client care plan. If a change in the client’s status occurs i.e. complications resulting from PN administration or frequent changes in PN orders, the plan of care is no longer considered established.

High-Alert Medication - medications that bear a heightened risk of causing significant patient harm when used in error as defined by the Institute for Safe Medication Practices (ISMP).

Independent Double-Check - the process where two clinicians separately check (alone and apart from each other, then compare results) each component of prescribing, dispensing and verifying the high-alert

medication for errors before it is administered to the client. The clinician checking has to form an independent judgment without cues from the clinician doing the initial work.

Parenteral Nutrition (PN) – administration of nutrients via venous route.

Peripheral – formulation appropriate for delivery via peripheral line.

Standardized Commercially Available PN – a commercially manufactured PN product.

Verification – a visual check that the correct medication, dose, rate and route is being administered according to the current prescribed medication order.

ROLES:

Graduate Nurses (GNs) – as assigned, GNs will administer and maintain PN with direct supervision until determined by an RN supervisor to be competent to practice autonomously.

Licensed Practical Nurses (LPNs) – LPNs identified by their manager in targeted practice settings will be certified in the LPN Additional Competency: Parenteral Nutrition – Adult Administration and Maintenance with an Established Plan of Care, to provide care independently as assigned, for clients who are less complex, more predictable and at lower risk for negative outcomes. LPN practice is limited to clients receiving PN through a peripheral IV or PICC line (or another type of CVC if certified). If a change is required in the established plan of care, the LPN will consult with a RN, RPN or physician and work collaboratively to establish a new plan of care. **Prerequisite:** LPN must have completed Sask. Polytechnic IV Therapy/Blood & Blood Products Completer Course or equivalent.

Registered Nurses (RNs) – as assigned, RNs will administer and maintain PN. If a change is required to a plan of care within an LPN's assignment, an RN will provide consultation as needed and work collaboratively with the LPN until a new plan of care is established.

Registered Psychiatric Nurse (RPN) who has the knowledge and skill may administer PN by peripheral IV or PICC if certified in care and use of PICCs.

1. PURPOSE

- 1.1 To ensure clients receive adequate nutritional support
- 1.2 To safely administer parenteral nutrition
- 1.3 To decrease the risk of infection, refeeding syndrome and other complications of PN administration

2. POLICY

- 2.1 The LPN certified in this Additional Competency will have first completed the following learning modules/activities prior to care for client on PN with established plan of care
 - 2.1.1 Complete the required learning module and quiz (teaching and learning methods may vary e.g. classroom and/or self- study using paper module or on line).

- 2.1.2 Complete a skills checklist with a certified RN or certified LPN during simulation or during first PN care to ensure safety checks are followed appropriately.
- 2.1.3 Provide documentation of learning module quiz and skills checklist to educator/supervisor
- 2.2 The Dietitian must be consulted to determine the client’s nutritional status and appropriate feeding route.
- 2.3 The most responsible physician will determine the need for PN and provide clinical supervision and monitoring of administration.

Note: At RUH, a consult to the Nutrition Support Service (NSS) is required.

- 2.4 PN orders will be written by the most responsible physician or designate using the “3-in1 Adult Parenteral Nutrition” order form (See Appendix A), “2-in-1 Adult Parenteral Nutrition” order form (See Appendix B) or the “Standardized Commercially Available Adult 3-in-1 Parenteral Nutrition” order form (See Appendix C). Orders must be written daily for new PN starts. Orders can remain active for several days once the client’s electrolyte and fluid status stabilizes. See the bottom of the order set for the date the next PN re-order is required.

Note: If the client is stabilized on PN, orders may be written once weekly on Thursdays.

Note: At RUH, NSS or (when unavailable) the unit Resident will write the orders.

- 2.5 All PN formulations will be independently double checked as per the Regional High Alert Medication Policy.
- 2.6 PN Orders
 - 2.6.1 Pharmacy must receive early notification (by 1100 hrs) for new PN starts.
 - 2.6.2 Orders must be written and scanned to pharmacy before 1100 hours. Pharmacy may not have the resources to compound late orders. Alternatives to compounded PN include standardized commercially available PN or dextrose containing IV fluids.
 - 2.6.3 The Dietitian may choose to use the standardized commercially available PN formulation when a client has a new indication for PN and it is after the 1100 hours deadline. In this case, the client may be transitioned to compounded PN the following day.
- 2.7 For all new PN clients an appropriate infusion line for administration must be in place. If the infusion line is peripheral, a 20 or 22 gauge IV should be inserted into the largest, straightest vein available. This will allow blood flow around the catheter therefore minimizing irritation to the vein.
- 2.8 When using a multiple lumen Central Venous Catheter (CVC), one lumen should be dedicated to PN administration.

Note: PN exceeding 10% dextrose or an osmolarity of greater than 900 mOsm/L is administered through a central venous catheter.

Note: No other IV solutions, medications, or blood products may be administered using the PN line (primary or secondary) or lumen. Replacement fluids must be infused separately.

Note: No blood withdrawal or CVP monitoring should be done using the PN lumen.

2.9 No additions may be made to the PN bag.

2.10 Administer PN at the ordered rate. The flow rate must not be increased to catch up if the infusion falls behind the ordered rate.

2.11 Peripheral PN may be infused through a central line until the current bag is completed then the formula must be changed to a central formulation.

Note: PN formulated for central use CANNOT be infused through a peripheral line.

2.12 The most responsible physician or designate will be notified if there are any signs of inflammation or discharge at the CVC or IV entry site.

2.13 Clients requiring community PN on discharge may be referred to and supported by the Provincial PN program.

3. PROCEDURE

3.1 Pre-Administration

3.1.1 Supplies:

- PN solution and filter from Pharmacy
- Infusion pump tubing
- Alcohol 70% swab
- Infusion pump
- Clean gloves

Note: For 2-in-1 formulations, Pharmacy will send primary infusion pump tubing with inline 0.2 micron filter with the non-lipid bag.

For 3-in-1 formulations a 1.2 micron filter placed close to the patient is required.

3.1.2 Change the PN bag, tubing and filter every 24 hours. Keep the PN bag refrigerated until 30 minutes prior to hanging.

3.2 Administration

3.2.1 Check the PN order for the flow rate of the formulation.

3.2.1.1 Initiate PN therapy as ordered and do not adjust the rate of PN unless ordered.

Note: Initial PN therapy will be ordered to begin at a slow rate and will be gradually increased over 48-72 hours to the desired daily volume due to the relatively high dextrose load.

Note: *In certain circumstances, (i.e. insulin dependent diabetics or unconscious clients) sudden discontinuation of PN should be followed by Dextrose 10% for 2-4 hours at 50 mls/hr as ordered by physician.*

3.2.1.2 Do not increase or decrease the rate of PN abruptly. If suspended, call for orders.

Note: *PN containing lipid can be weaned over 2 hours if necessary.*

3.2.2 Compare the PN label to the client's name, the ordered formulation and electrolyte additions (2 –in-1, 3-in-1 or standardized commercially available formulation) against the practitioner's order. Check the expiry date and the bag for presence of particulate matter or creaming. PN should not be infused if visual changes or precipitates are apparent.

Note: *All PN formulations require an independent double check of the PN formulation against the practitioner order.*

3.2.3 Rotate or agitate the bag prior to hanging to keep the 3-in-1 formulation or standardized commercially available formulation from settling (the standardized commercially available formulation will be activated by pharmacy and will arrive on the nursing unit as a single bag).

3.2.4 Perform hand hygiene.

3.2.5 Prime the tubing and filter (refer to instructions on the filter package)

Note: *All 3-in-1 formulations require a 1.2 micron filter placed closest to the client (See Appendix D).*

Note: *For 2-in-1 formulations, the non-lipid bag is hung on the primary line using pump tubing with inline 0.2 micron filter sent by Pharmacy. The lipid bag is hung on the secondary line. Connect the provided 1.2 micron filter to the pump tubing closest to the client (See Appendix E).*

Note: *For administering only the non-lipid formulation, the non-lipid bag is hung on the primary line using pump tubing with inline 0.2 micron filter sent by Pharmacy (See Appendix F). A 1.2 micron filter is not needed.*

3.2.6 Turn off existing PN line.

3.2.7 If administering PN via a CVC with a clamp, clamp the catheter to prevent air embolism and/or blood loss.

3.2.8 Perform hand hygiene. Don clean gloves.

3.2.9 Needleless adaptor must be scrubbed for 15 seconds using Alcohol 70% swab and friction in a twisting motion prior to accessing (let dry).

3.2.10 Disconnect old tubing, avoiding contamination of the central venous or IV catheter.

3.2.11 Connect new tubing and filter.

- 3.2.12 Check all connections.
- 3.2.13 Unclamp catheter and/or tubing if clamp in place.
- 3.2.14 Remove gloves. Perform hand hygiene.
- 3.2.15 Program the infusion pump and start PN infusion as ordered.

Note: All PN formulations require an independent double check at the pump.

Note: Malnourished clients may have slower than normal rates of infusion to prevent refeeding syndrome (See Appendix G).

Note: Administering the lipid component of a 2-in-1 formulation at the incorrect rate may cause fat overload syndrome (See Appendix G).

3.3 Monitoring

- 3.3.1 Weigh the client once a week and document the weight in kilograms on appropriate record.
- 3.3.2 Maintain accurate intake and output records.
- 3.3.3 Monitor lab work according to the Adult Blood Work Monitoring for PN form (See Appendix H).
- 3.3.4 Document date and time of tubing, filter and bag change on appropriate record.

4. REFERENCES

American Society for Parenteral and Enteral Nutrition. (2016). *Parenteral Nutrition – New Recommendations for In-line Filters*. Retrieved January 6, 2017

from http://www.nutritioncare.org/News/General_News/Parenteral_Nutrition_%E2%80%93_New_Recommendations_for_In-line_Filters/

Ayers, P., et al. (2014). A.S.P.E.N. Parenteral Nutrition Safety Consensus Recommendations. *Journal of Parenteral and Enteral Nutrition*, 38(3), 296–333.

Ball, P. A. (2003). Intravenous in-line filters: filtering the evidence.

Boullata, J. I., Gilbert, K., Sacks, G., Labossiere, R. J., Crill, C., Goday, P., Kumpf, V.J., Mattox, T.W., Plogsted, S., & Holcombe, B. (2014). A.S.P.E.N. Clinical Guidelines: Parenteral Nutrition Ordering, Order Review, Compounding, Labeling, and Dispensing. *Journal of Parenteral and Enteral Nutrition*, 38(3), 334–377.

Correia, M. I. T. D., Guimaraes, J., de Mattos, C., Araujo Gurgel, K. C., & Cabral, E. B. (2004). Peripheral parenteral nutrition: an option for patients with an indication for short-term parenteral nutrition. *Nutricion Hospitalaria*, 1, 14–18.

Dietitians Association of Australia. (2015). Parenteral nutrition manual for adults in health care facilities.

Fresenius, K. (2015). SMOFlipid™ - Product monograph, lipid emulsion for intravenous nutrition. Retrieved from <https://www.fresenius-kabi.com/en-ca/documents/SMOFlipid-PM-En-2015.pdf>

Guenter, P., Ayers, P., Boullata, J. I., Gura, K. M., Holcombe, B., & Sacks, G. S. (2017). Parenteral nutrition errors and potential errors reported over the past 10 years. *Nutrition in Clinical Practice*, 32(6), 826–830.

Hojsak, I., & Kolaček, S. (2014). Fat overload syndrome after the rapid infusion of SMOFlipid™ emulsion. *Journal of Parenteral & Enteral Nutrition*, 38(1), 119–121.

Inayet, N., & Neild, P. (2015). Parenteral nutrition. *The Journal of the Royal College of Physicians of Edinburgh*, 45, 45–48.

Institute for Safe Medication Practices (ISMP). (2012). ISMP's list of high-alert Medications. Retrieved from <http://www.ismp.org/tools/highalertmedications.pdf>

National Collaborating Centre for Acute Care. (2006). Nutrition support in adults: Oral nutrition support, enteral tube feeding and parenteral nutrition. National Collaborating Centre for Acute Care, London. Retrieved from www.rcseng.ac.uk

O'Grady, N.P., Alexander, M., Burns, L, et al. (2011). *Guidelines for the Prevention of Intravascular Catheter-Related Infections*, 52(9). Center for Disease Control.

Phillips, L. D. (2010). *Manual of IV therapeutics* (5th ed.). Philadelphia, PA: F. A. Davis Company.

Pittiruti, M., Hamilton, H., Biffi, R., Macfie, J., & Pertkiewicz, M. (2009). ESPEN Guidelines on Parenteral Nutrition: Central Venous Catheters (access, care, diagnosis and therapy of complications). *Clinical Nutrition*, 28(4), 365–377.

Sacks, G. S., Rough, S., & Kudsk, K. A. (2009). Frequency and severity of harm of medication errors related to the parenteral nutrition process in a large university teaching hospital. *Pharmacotherapy*, 29(8), 966–974.



Ukleja, A. et al. (2010). Standards for Nutrition Support: Adult Hospitalized Patients. *Nutrition in Clinical Practice*, 25(4), 403-414.

Worthington, P. H., & Gilbert, K. A. (2012). Parenteral nutrition. Risks, complications, and management. *Infusion Nurses Society*, 35(1), 52–64.


Worthington, P. et al (2017). When is parenteral nutrition appropriate? *Journal of Parenteral and Enteral Nutrition*, 41(3), 324–377.

Yantis, M A., & Velandar, R. (2009). How to recognize and respond to refeeding syndrome. *Nursing 2009 Critical Care*, 4(3), 14-20.

Appendix A


		PATIENT IDENTIFICATION <input type="checkbox"/> RUH <input type="checkbox"/> SCH <input type="checkbox"/> SPH <input type="checkbox"/> OTHER: _____				
3-in-1 ADULT PARENTERAL NUTRITION		ACTION <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 25%;">MAR</td> <td style="width: 25%;">TOP</td> <td style="width: 25%;">REC</td> <td style="width: 25%;">RN</td> </tr> </table>	MAR	TOP	REC	RN
MAR	TOP	REC	RN			
<input checked="" type="checkbox"/> Consult the Nutrition Support Service at RUH. Consult the dietitian at SPH or SCH. <input checked="" type="checkbox"/> NEW orders MUST be RECEIVED in Pharmacy NO LATER than 1100 hours or the formulation from the previous day will be supplied. <input checked="" type="checkbox"/> Start Adult Blood work Monitoring for Parenteral Nutrition (Form #101758)		Route of Administration <input type="checkbox"/> CVC <input type="checkbox"/> Peripheral IV				
Primary diagnosis: _____ Indication for parenteral nutrition: _____		Height _____ cm Dosing Weight _____ kg				
Select the base: <input type="checkbox"/> 3-in-1 Standard Central Amino Acids (Travasol) 50 g/L Dextrose 165 g/L Lipid Emulsion (SMOF) 30 g/L Total Calories 1081 Kcal/L <input type="checkbox"/> 3-in-1 Intermediate Central Amino Acids (Travasol) 55 g/L Dextrose 135 g/L Lipid Emulsion (SMOF) 25 g/L Total Calories 937 Kcal/L <input type="checkbox"/> 3-in-1 Stress Central Amino Acids (Travasol) 63 g/L Dextrose 100 g/L Lipid Emulsion (SMOF) 30 g/L Total Calories 892 Kcal/L <input type="checkbox"/> 3-in-1 Peripheral Amino Acids (Travasol) 30 g/L Dextrose 70 g/L Lipid Emulsion (SMOF) 35 g/L Total Calories 708 Kcal/L	Order daily additions: sodium _____ mEq/day potassium _____ mEq/day calcium _____ mEq/day magnesium _____ mEq/day phosphate _____ mmol/day insulin HumuLIN R (Regular) _____ units/day <input checked="" type="checkbox"/> vitamin K 150 mcg/day <input checked="" type="checkbox"/> multivitamins 10 mL/day <input checked="" type="checkbox"/> trace elements (Micro+6) 1 mL/day <input checked="" type="checkbox"/> maximize chloride unless alternative is selected <input type="checkbox"/> equal chloride to acetate (Approximate) <input type="checkbox"/> maximize acetate					
Optional Trace Elements (Order ONLY if required): <input type="checkbox"/> Do not use Micro+6 trace elements. Use Micro+4 Regular 5 mL/day <input type="checkbox"/> Extra chromium _____ mcg/day (Max 20 mcg/day) <input type="checkbox"/> Extra copper _____ mg/day (Max 2.4 mg/day) <input type="checkbox"/> Extra selenium _____ mcg/day (Max 200 mcg/day) <input type="checkbox"/> Extra zinc _____ mg/day (Max 15 mg/day)						
Order rate of administration: Requires a 1.2 micron filter connected to the end of the primary tubing. <input type="checkbox"/> Continuous infusion _____ mL/h for 24 hours <input type="checkbox"/> Cyclic infusion: _____ mL/h for _____ hours (from _____ h to _____ h) then _____ mL/h for _____ hours (from _____ h to _____ h) then _____ mL/h for _____ hours (from _____ h to _____ h)						
A daily order is required for the first 7 days. PN can be reordered once a week on Thursdays for stable patients.						
Order Date _____ <input type="checkbox"/> Review labs & reorder tomorrow <input type="checkbox"/> Review labs & reorder on _____		<input type="checkbox"/> Review labs & reorder every Thursday (default)				
Dietitian Name and Pager _____		First day of PN was _____				
<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 33%; height: 20px;"> </td> <td style="width: 33%; height: 20px;"> </td> <td style="width: 33%; height: 20px;"> </td> </tr> </table>						
_____ PRACTITIONER PRINTED NAME	_____ PRACTITIONER SIGNATURE	_____ DATE/TIME				
Form #100020	10/17	Category: Orders				
		TPN3				
		Page 1 of 1				

Appendix C

	PATIENT IDENTIFICATION <input type="checkbox"/> RUH <input type="checkbox"/> SCH <input type="checkbox"/> SPH <input type="checkbox"/> OTHER: _____					
Standardized Commercially Available Adult 3-in-1 Parenteral Nutrition		ACTION <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 25%; text-align: center;">MAR</td> <td style="width: 25%; text-align: center;">LOP</td> <td style="width: 25%; text-align: center;">REC</td> <td style="width: 25%; text-align: center;">RN</td> </tr> </table>	MAR	LOP	REC	RN
MAR	LOP	REC	RN			
<input checked="" type="checkbox"/> Consult the Nutrition Support Service at RUH. Consult the dietitian at SPH or SCH. <input checked="" type="checkbox"/> Use for a new start parenteral nutrition order written between 1100h and 1700h <input checked="" type="checkbox"/> Discontinue this order when new orders are written <input checked="" type="checkbox"/> Any change requires completion of a new order set (do not alter or edit completed orders) <input checked="" type="checkbox"/> Start Adult Blood work Monitoring for Parenteral Nutrition (Form #101758) <input checked="" type="checkbox"/> No additions will be made to the commercial base		Route of Administration <input type="checkbox"/> CVC or PICC <input type="checkbox"/> Peripheral IV				
Primary diagnosis: _____ Indication for parenteral nutrition: _____		Height _____ cm Dosing Weight _____ kg				
<input type="checkbox"/> Olime! 5.7% E Central Amino Acids 58.9 g/L Dextrose 110 g/L Lipid Emulsion (Clinoleic) 40 g/L Total Calories 1070 Kcal/L Administer through a central line (CVC or PICC)	Electrolyte Contents: sodium 35 mEq/L potassium 30 mEq/L calcium 7 mEq/L magnesium 8 mEq/L phosphate 15 mmol/L					
<input type="checkbox"/> PeriOlime! 2.5% E Peripheral Amino Acids 25.3 g/L Dextrose 75 g/L Lipid Emulsion (Clinoleic) 30 g/L Total Calories 700 Kcal/L Administer through a peripheral line.	Electrolyte Contents: sodium 21 mEq/L potassium 16 mEq/L calcium 4 mEq/L magnesium 4.4 mEq/L phosphate 8.5 mmol/L					
Order rate of administration: All 3-in-1 parenteral nutrition requires a 1.2 micron filter placed closest to the patient <input type="checkbox"/> Continuous infusion _____ mL/h for 24 hours <input type="checkbox"/> Cyclic infusion: _____ mL/h for _____ hours (from _____ h to _____ h) then _____ mL/h for _____ hours (from _____ h to _____ h) then _____ mL/h for _____ hours (from _____ h to _____ h)						
This order is only active for 1 day						
Dietitian Name and Pager _____		Date/Time _____				
<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 33%; height: 20px;"></td> <td style="width: 33%;"></td> <td style="width: 33%;"></td> </tr> </table>						
PRACTITIONER PRINTED NAME _____	PRACTITIONER SIGNATURE _____	DATE/TIME _____				
<small>Notice of confidentiality: Contains information that is time sensitive or confidential. Use, disclosure, copying or communication of the contents is prohibited. If you have received in error, notify the SHR Pharmacy Manager, Operations (306-666-6696).</small>						
Form #104114	07/16	Category: Orders TPNCOM				
Page 1 of 1						


Appendix D

**Instructions for the Administration of 3 in 1 PN
(compounded formulation or the standardized commercially available formulation)**




Appendix E

Special Instructions for the Administration of 2 in 1 PN



Appendix F

Special Instructions for the Administration of Non-Lipid PN Only



Appendix G

Fat Overload Syndrome

Fat overload syndrome is a complication due to parenteral lipids being infused faster than indicated and can cause the following:

Hyperlipidemia	Fever	Fat Infiltration
Hepatomegaly (may have jaundice)	Splenomegaly	Anemia
Leukopenia	Thrombocytopenia	Coagulation Disorders
Hemolysis	Reticulocytosis	Abnormal Liver Function Test
Coma		

Refeeding Syndrome

Refeeding syndrome is a fluid and electrolyte imbalance after nutritional support such as intravenous fluid, enteral tube feeding or oral intake has started on a malnourished client. This will cause:

- Hypokalemia
- Hypophosphatemia
- Hypomagnesemia

This in turn can cause:

CNS: Weakness Altered mental state Seizure Paresthesia Paralysis Tetany	GI: Nausea and vomiting Anorexia Constipation/diarrhea Abdominal pain Vitamin deficiency Low albumin/prealbumin	Hematologic: Bleeding Infection Anemia Thrombocytopenia Platelet dysfunction
CVS: Hypotension Dysrhythmias	GU: Edema Elevated BUN & Creatinine	Metabolic: Metabolic alkalosis/acidosis
Respiratory: Hypoxia	Musculoskeletal: Myalgia	

Appendix H



SASKATOON HEALTH REGION
Saskatoon, Saskatchewan

RUH SCH SPH Other _____

Patient Label

NAME: _____

HSN: _____

D.O.B.: _____

ADULT BLOODWORK MONITORING FOR PN

Page 1 of 2

	WEEK ONE					WEEK TWO		
DATE								
DAY OF PN	1	2	3	4	6	Mon	Weds	Fri
PANEL A	X			X*		X		
PANEL B		X	X				X	
PANEL C					X			X

	WEEK THREE		WEEK FOUR		WEEK FIVE	
DATE						
DAY OF PN	Mon	Thurs	Mon	Thurs	Mon	Thurs
PANEL A	X				X	
PANEL B		X		X		X
PANEL D			X			

PANEL A includes:	PANEL B includes:	PANEL C includes:	PANEL D includes:
Hematology: CBC, INR Chemistry: Lytes ⁷ Prealbumin C-Reactive Protein Phosphate Magnesium Ionized Calcium Triglycerides AST, ALT, ALP, GGT Direct & Total Bilirubin * On Day 4 of week one do not order Prealbumin and C-reactive protein	Chemistry: Lytes ⁷ Phosphate Magnesium	Chemistry: Lytes ⁷ Phosphate Magnesium Ionized Calcium Triglycerides	Panel A <u>plus</u> Zinc Selenium Iron TIBC Ferritin *Drawn every 4 weeks starting on the Monday of Week 4
Physician's Signature: _____			