

**Trained Observer Checklist for Personal Protective Equipment (PPE)  
Doffing – High Risk<sup>1</sup>**

Done ✓	
	Review page 1 of HCW D <b>OFF</b> ING PPE Procedure *HIGH RISK
	Trained observer is wearing shoe covers, gown, face shield, gloves
	Trained observer is ready to monitor the steps of the Checklist for PPE Doffing and reading aloud each step of the procedure
	Prior to taking <b>off</b> PPE, the trained observer will remind the HCW to avoid touching eyes, nose, mouth and to perform each step slowly
	1. Take <b>off</b> outside gloves inside client room
	2. Trained observer visually inspects HCW PPE
	3. Take <b>off</b> gown
	4. Take <b>off</b> gloves
	5. Hand hygiene
	6. Put <b>on</b> gloves
	7. Take <b>off</b> shoe cover
	8. Take <b>off</b> gloves
	9. Hand hygiene
	10. Put <b>on</b> gloves
	11. Take <b>off</b> face shield
	12. Take <b>off</b> gloves
	13. Hand hygiene
	14. Put <b>on</b> gloves
	15. Take <b>off</b> surgical hood
	16. Take <b>off</b> gloves
	17. Hand hygiene
	<b>18. Remove N95 respirator – remind HCW not to touch face after removal until hand hygiene is performed</b>
	19. Hand hygiene
	20. Put <b>on</b> gloves
	21. Clean shoes
	22. Take <b>off</b> gloves
	23. Hand hygiene
	24. Trained observer to pick up any items that fell to floor
	25. Trained observer to take <b>off</b> PPE using the Trained Observer DOFFING PPE

Date/Time \_\_\_\_\_

Healthcare Worker \_\_\_\_\_

Trained Observer \_\_\_\_\_

<sup>1</sup>High risk symptoms: Bleeding or uncontrolled diarrhea or uncontrolled vomiting

High risk procedures: Aerosol generating medical procedures (AGMPs), CPR, central line insertion, procedures that result in copious amounts of body fluid contamination